

LUTTINGER SURGERY AND INTERESTING SYMPLECTIC 4-MANIFOLDS WITH SMALL EULER CHARACTERISTIC

SCOTT BALDRIDGE AND PAUL KIRK

ABSTRACT. In this article we construct a minimal symplectic 4-manifold R that has small Euler characteristic ($e(R) = 8$) and two essential Lagrangian tori with nice properties. These properties make R particularly suitable for constructing interesting examples of symplectic manifolds with small Euler characteristic. In particular, we construct an exotic symplectic $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$, the smallest known minimal symplectic 4-manifold with $\pi_1 = \mathbb{Z}$, the smallest known minimal symplectic 4-manifolds with $\pi_1 = \mathbb{Z}/a \oplus \mathbb{Z}/b$ for all $a, b \in \mathbb{Z}$, and the smallest known minimal symplectic 4-manifold with $\pi_1 = \mathbb{Z}^3$. We use the $\pi_1 = \mathbb{Z}$ example to derive a significantly better upper bound on the minimal Euler characteristic of all symplectic 4-manifolds with a prescribed fundamental group.

1. INTRODUCTION

Combining the construction of taking symplectic sums along a genus 2 surface in $(T^2 \times S^2) \# 4\overline{\mathbb{CP}}^2$ (which we learned from a recent article of article of Akhmedov [2] and also an article of Ozbaggi and Stipsitz [18]), with the method of (symplectic) Luttinger surgery ([15, 1]) along Lagrangian tori, in this article we prove the following theorem (see Theorem 6).

Theorem. *There exists a minimal symplectic 4-manifold R containing a pair of homologically essential Lagrangian tori T_1, T_2 . The manifold R satisfies*

- *The Euler characteristic $e(R) = 8$, the signature $\sigma(R) = -4$,*
- *the meridians to T_1 and T_2 are nullhomotopic in $R - (T_1 \cup T_2)$,*
- *$\pi_1(R) = \mathbb{Z}^2$ and the map $\pi_1(R - (T_1 \cup T_2)) \rightarrow \pi_1(R)$ is an isomorphism,*
- *the inclusions induce the homomorphisms $\mathbb{Z}^2 = \pi_1(T_1) \rightarrow \pi_1(R), (s, t) \mapsto (s, 0)$ and $\mathbb{Z}^2 = \pi_1(T_2) \rightarrow \pi_1(R), (s, t) \mapsto (0, t)$.*

Combined with techniques such as Gompf's symplectic sum [12] and Luttinger surgery [1] we use R to construct interesting examples of minimal symplectic 4-manifolds with prescribed fundamental group and small Euler characteristic. In particular, we construct:

- (1) A symplectic manifold M with $\pi_1(M) = \mathbb{Z}$, $e(M) = 8$, $\sigma(M) = -4$. This manifold contains a symplectic torus T with trivial normal bundle which can be used as a smaller replacement to the elliptic surface $E(1)$ to kill elements in fundamental groups of symplectic 4-manifolds (Theorem 7).
- (2) For each group $G = \mathbb{Z}/a \oplus \mathbb{Z}/b$, with a, b arbitrary integers, an infinite family of minimal symplectic 4-manifolds with fundamental group G , Euler characteristic 8 and signature -4 (Theorem 9). The case $a = b = 1$ yields minimal symplectic 4-manifolds homeomorphic to $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$.
- (3) For each group G presented with g generators and r relations, a minimal symplectic 4-manifold with fundamental group G , Euler characteristic $12 + 8(g + r)$ and signature $-8 - 4(g + r)$. This significantly improves the main result of [6].

The authors would like to thank A. Akhmedov, R. Fintushel, C. Judge, C. Livingston, and J. Yazinsky for helpful discussions.

Date: January 23, 2007.

2000 Mathematics Subject Classification. Primary 57R17; Secondary 57M05, 54D05.

Key words and phrases. Symplectic topology, fundamental group, 4-manifold.

The first author gratefully acknowledges support from the NSF grant DMS-0507857. The second author gratefully acknowledges support from the NSF grant DMS-0604310.

2. SOME TOPOLOGY

Consider the union of three symplectic surfaces $(T^2 \times \{s_1\}) \cup (\{r\} \times S^2) \cup (T^2 \times \{s_2\})$ in the symplectic 4-manifold $T^2 \times S^2$ (with the product symplectic form). Resolving the three double points symplectically ([12]) yields a genus 2 symplectic surface F with self-intersection 4. Blowing up $T^2 \times S^2$ along four points on F and taking the proper transform yields a genus 2 surface (which we continue to call F) in

$$W = (T^2 \times S^2) \# 4\overline{\mathbb{CP}}^2.$$

The symplectic surface $F \subset W$ has the easily verified properties:

- (1) Based loops a_1, b_1, a_2, b_2 on F representing the standard symplectic generators can be chosen so that $\pi_1(F) = \langle a_1, b_1, a_2, b_2 \mid [a_1, b_1][a_2, b_2] = 1 \rangle$ in such a way that $\pi_1(F)$ surjects to $\pi_1(W) = \mathbb{Z}a \oplus \mathbb{Z}b$ by the assignment
$$a_1 \mapsto a, b_1 \mapsto b, a_2 \mapsto a^{-1}, b_2 \mapsto b^{-1}.$$
- (2) F intersects each of the four -1 exceptional spheres transversely once.
- (3) F has a trivial normal bundle.

Fix a trivialization of the normal bundle of F , and hence an identification of the boundary of a tubular neighborhood of F with $F \times S^1$. The meridian of F is the curve $\{p\} \times S^1 = \mu_F \subset W - F$ which is the boundary of a small normal disc to F . Up to a (free) homotopy, we may assume that μ_F lies on one of the exceptional spheres, and hence μ_F is nullhomotopic in $W - F$. In particular, since F is connected, every homotopy that intersects F can be replaced by a homotopy that misses F . Since every loop in W can be pushed off F it follows that the inclusion $W - F \subset W$ induces an isomorphism on fundamental groups. Hence $\pi_1(W - F) = \mathbb{Z}a \oplus \mathbb{Z}b$, and the push off of F into $W - F$ using the trivialization again induces the homomorphism $a_1 \mapsto a, b_1 \mapsto b, a_2 \mapsto a^{-1}, b_2 \mapsto b^{-1}$.

(Matsumoto [16] describes a Lefschetz fibration $f : W \rightarrow S^2$ with generic fiber F and with eight singular fibers; the monodromy given by the relation $(D_1 D_2 D_3 D_4)^2 = 1$ in the mapping class group of F , where D_i is the positive Dehn twist about the curve C_i of [13, pg 325].)

Lemma 1. *Let R be any 4-manifold containing a genus 2 surface G with trivialized normal bundle. Let $\phi : F \rightarrow G$ be a diffeomorphism, and set $g_i = \phi_*(a_i), h_i = \phi_*(b_i)$ in $\pi_1(R)$. Given a map $\tau : F \rightarrow S^1$, let $\tilde{\phi} : F \times S^1 \rightarrow G \times S^1$ the diffeomorphism given by $\tilde{\phi}(a, s) = (\phi(a), \tau(a) \cdot s)$. Then the sum*

$$S = R \#_{F,G} W = (R - \text{nb}d(G)) \cup_{\tilde{\phi}} (W - \text{nb}d(F))$$

has fundamental group

$$\pi_1(S) = \pi_1(R) / N(g_2 g_1, h_2 h_1, [g_1, h_1])$$

where $N(g_1 g_2, h_1 h_2, [g_1, h_1])$ denotes the normal subgroup generated by $g_1 g_2$, $h_1 h_2$, and the commutator $[g_1, h_1]$.

Proof. The homomorphism $\pi_1(R - G) \rightarrow \pi_1(R)$ is a surjection since every loop can be pushed off G , and the kernel is normally generated by the meridian $\mu_G = \tilde{\phi}(\mu_F)$ because any homotopy can be made transverse to G and G is connected.

The Seifert-Van Kampen theorem implies that $\pi_1(S)$ is the quotient of the free product $\pi_1(R - G) * \pi_1(W - F) = \pi_1(R - G) * (\mathbb{Z}a \oplus \mathbb{Z}b)$ by the normal subgroup generated by $\tilde{\phi}(\mu_F) \mu_G^{-1}, g_1 a^{-1}, h_1 b^{-1}, g_2 a, h_2 b$. Since $\mu_F = 1$, this can also be described as the quotient of $\pi_1(R) * (\mathbb{Z}a \oplus \mathbb{Z}b)$ by the normal subgroup generated by $g_1 a^{-1}, h_1 b^{-1}, g_2 a, h_2 b$. We can then eliminate the generators a and b and conclude that $\pi_1(S)$ is the quotient of $\pi_1(R)$ by the normal subgroup generated by $g_2 g_1, h_2 h_1$, and the commutator $[g_1, h_1]$. $\square$

We next construct a useful building block for our subsequent constructions. We will postpone the discussion of symplectic structures until the next section. Our emphasis now is on a careful calculation of fundamental groups.

We begin with the statement of a well-known lemma which computes the fundamental groups of generalized mapping tori. We omit the standard proof.

Lemma 2. *Let X be a finite connected CW complex with base point x_0 , $i : Y \subset X$ the inclusion of a connected subcomplex containing x_0 , and $f : Y \rightarrow X$ a continuous base point preserving map. Suppose $\pi_1(X, x_0)$ is generated by loops $x_1, \dots, x_g$, with a presentation $\langle x_1, \dots, x_g \mid w_1, \dots, w_r \rangle$, and $\pi_1(Y, x_0)$ is generated by loops $\gamma_1, \dots, \gamma_n$.*

Let $Z = X \times [0, 1] / \sim$ where $(y, 0) \sim (f(y), 1)$ for $y \in Y$. Give Z the base point $z_0 = (x_0, 0)$ and denote by $x_i \in \pi_1(Z, z_0)$ again the images of the loops x_i under the map $X = X \times \{0\} \subset X \times [0, 1] \rightarrow Z$. Let $t \in \pi_1(Z, z_0)$ denote the loop $u \mapsto (x_0, u) \subset X \times [0, 1] \rightarrow Z$.

Then $\pi_1(Z, z_0)$ has the presentation

$$\pi_1(Z, z_0) = \langle t, x_1, \dots, x_g \mid w_1, \dots, w_r, ti_*(\gamma_1)t^{-1} = f_*(\gamma_1), \dots, ti_*(\gamma_n)t^{-1} = f_*(\gamma_n) \rangle.$$

□

Let H be a once punctured torus. Choose a base point $h \in H$ on the boundary of H and denote by $x, y \in \pi_1(H, h)$ the two standard symplectic free generators of the fundamental group. Let $D : H \rightarrow H$ denote the Dehn twist along a curve parallel to x . Let $K \subset H$ denote a loop parallel to y and L a loop parallel to x , as illustrated in the following figure.




FIGURE 1. The surface H .

Let Z denote the mapping torus of $D : H \rightarrow H$, i.e.

$$Z = H \times [0, 1] / \sim \text{ where } (x, 0) \sim (D(x), 1).$$

Thus Z is a fiber bundle over $S^1 = [0, 1] / (0 \sim 1)$ with fibers H . Let $C = Z \times S^1$. For convenience we think of the second coordinate as $[0, 1] / (0 \sim 1)$. Hence we give C coordinates $(a, u, v) \in H \times [0, 1] \times [0, 1]$. We have loops $x = x \times (0, 0)$, $y = y \times (0, 0)$, $t = \{h\} \times I \times \{0\}$, and $s = \{h\} \times \{0\} \times I$ all based at $h = (h, 0, 0)$ (note that D fixes h , so that t and s are loops). Applying Lemma 2 gives:

$$\pi_1(C) = \langle x, y, t \mid txt^{-1} = x, tyt^{-1} = yx \rangle \oplus \mathbb{Z}s = \langle x, y, t \mid [t, x], [y^{-1}, t]x^{-1} \rangle \oplus \mathbb{Z}s.$$

There are two interesting tori in C . One is the torus $T_1 = K \times \{0\} \times I / \sim$, the other is the torus $T_2 = L \times I \times \{0\} / \sim$. Notice that the definition of T_2 makes sense since L misses the support of the Dehn twist D .

We connect T_1 and T_2 to the base point $(h, 0, 0)$ as follows. For T_1 , follow the path in $H \times \{(0, 0)\}$ starting at $(h, 0, 0)$ illustrated in Figure 1 to the intersection of the loops labeled x and y , then follow x in the reverse direction a short way until you reach $K = K \times \{(0, 0)\} \subset T_1$. For T_2 follow the path to the intersection of x and y , then follow y backwards until you hit $L = L \times \{(0, 0)\} \subset T_2$. Based this way, the two generators of $\pi_1(T_1)$ map to the classes y and s in $\pi_1(C)$, and the two generators of $\pi_1(T_2)$ map to the classes x and t in $\pi_1(C)$.

We are interested in the fundamental group of their complement and corresponding meridian circles. We begin with the calculation of $\pi_1(C - T_1)$. Notice that (identifying K with $K \times \{0\}$) $C - T_1 = (Z - K) \times S^1$, and hence $\pi_1(C - T_1) = \pi_1(Z - K) \oplus \mathbb{Z}s$. Moreover the meridian μ_1 of T_1 (the homotopy class of the boundary of a small 2-disk transverse in C to T_1) is represented by a loop in $Z - K$.

Since $Z - K$ is constructed as the quotient of $H \times [0, 1]$ where $(x, 0) \sim (D(x), 1)$ for $x \in H - K$, and $\pi_1(H - K)$ is generated by y and $c = [x, y] = \partial H$, it follows from Lemma 2 that

$$\pi_1(C - T_1) = \langle x, y, t \mid [y^{-1}, t]x^{-1}, [t, [x, y]] \rangle \oplus \mathbb{Z}s$$

and the map $\pi_1(C - T_1) \rightarrow \pi_1(C)$ takes x to x , y to y , t to t , and s to s . The meridian μ_1 is represented by $[t, x]$.

A similar argument computes $\pi_1(C - T_2)$: this time $C - T_2 = Z \times [0, 1]/\sim$, where $(z, 0) \sim (z, 1)$ when $z \in Z - T_2$. Thus the generator s only commutes with those elements of $\pi_1(Z)$ which are represented by loops that miss T_2 . Lemma 2 implies

$$\pi_1(C - T_2) = \langle x, y, t, s \mid [t, x], [y^{-1}, t]x^{-1}, [s, x], [s, t] \rangle$$

(with the obvious morphism to $\pi_1(C)$) and the meridian μ_2 is represented by $[s, y]$.

Finally, note that the boundary ∂C is a 3-torus, with $\pi_1(\partial C, h) = \mathbb{Z}c \oplus \mathbb{Z}t \oplus \mathbb{Z}s$ with $c = [x, y]$.

Remark. The only subtle parts of these calculations are the computations of μ_1 and μ_2 . However, an alternative and simpler calculation which is quite sufficient for our purposes is the following. Notice that the torus T_1 intersects T_2 transversally in one point, say p . Choose a pair of loops on T_2 which miss p and are freely homotopic to x and t . Connect these loops to the base point of C in $C - T_1$, and call the result x' and t' . Then obviously $[t', x']$ is a meridian for T_1 . Moreover, t' is conjugate to t and x' is conjugate to x . To anticipate what follows, notice that if $C - T_1$ is modified in some way so that x becomes nullhomotopic, then x' also becomes nullhomotopic and hence also $[t', x']$. The same argument shows that a meridian for T_2 is of the form $[s', y']$ for some curves s' conjugate to s and y' conjugate to y .

3. THE SYMPLECTIC MANIFOLD R

Let G denote another surface of genus 2, with base point g . Fix based loops x_1, y_1, x_2, y_2 representing a symplectic basis of $\pi_1(G)$. Let $D_1 : G \rightarrow G$ denote the Dehn twist about a curve parallel to x_1 , and let $D_2 : G \rightarrow G$ denote the Dehn twist about a curve parallel to y_2 . Let K be a curve parallel to y_1 and L a curve parallel to y_2 . These curves are illustrated in the following figure.




FIGURE 2. The surface G .

Denote by $\phi : G \rightarrow G$ the composite $\phi = D_2 \circ D_1$. The mapping torus of ϕ is a 3-manifold

$$Y = G \times [0, 1]/\sim, \text{ where } (x, 0) \sim (\phi(x), 1) \text{ for } x \in G.$$

The projection $Y \times [0, 1] \rightarrow [0, 1]$ determines a fibration $p : Y \rightarrow S^1$ of Y over the circle. Let $B = Y \times S^1$.

Notice that B is the union along their boundary T^3 of two copies of the manifold C constructed in the previous section, but with a reversal of the roles of x and y in the second copy (to the right in the figure). The Seifert-Van Kampen theorem coupled with the calculation of the previous section give

$$\pi_1(B) = \langle x_1, y_1, x_2, y_2, s, t \mid [x_1, y_1][x_2, y_2], [t, x_1], [y_1^{-1}, t]x_1^{-1}, [x_2^{-1}, t]y_2^{-1}, [t, y_2] \rangle \oplus \mathbb{Z}s.$$

As before, we view B as a quotient of $G \times [0, 1] \times [0, 1]$. We then have three disjoint surfaces in B : a pair of tori

$$T_1 = K \times \{0\} \times I / \sim$$

$$T_2 = L \times I \times \{0\} / \sim$$

and a genus two surface

$$G = G \times \{\frac{1}{2}\} \times \{\frac{1}{2}\} / \sim.$$

The tori T_1 and T_2 can be connected to the base point $(g, 0, 0)$ by the chosen path in $G \times \{(0, 0)\}$, and G can be connected to the base point using the path (g, u, u) , $u \in [0, \frac{1}{2}]$. The T_i correspond to the tori identified in the previous section (again with an appropriate reversal of x and y in the second copy of C), and so the Seifert-Van Kampen theorem can be used to compute $\pi_1(B - (T_1 \cup T_2))$ and the two corresponding meridians μ_1, μ_2 .

We summarize the resulting fundamental group calculations in the next proposition.

Proposition 3. *For the manifold B constructed above and the three surfaces T_1, T_2 and G we have*

$$\pi_1(B) = \langle x_1, y_1, x_2, y_2, t \mid [x_1, y_1][x_2, y_2], [t, x_1], [y_1^{-1}, t]x_1^{-1}, [x_2^{-1}, t]y_2^{-1}, [t, y_2] \rangle \oplus \mathbb{Z}s,$$

$\pi_1(G)$ is generated by x_1, y_1, x_2 , and y_2 , $\pi_1(T_1)$ is generated by y_1 and s , $\pi_1(T_2)$ is generated by y_2 and t .

Moreover, $\pi_1(B - (T_1 \cup T_2))$ is generated by x_1, y_1, x_2, y_2, t and s subject to the relations

$$[x_1, y_1][x_2, y_2] = [y_1^{-1}, t]x_1^{-1} = [t, [x_1, y_1]] = [x_2^{-1}, t]y_2^{-1} = [t, y_2] = [s, x_1] = [s, y_1] = [s, t] = [s, y_2] = 1.$$

In this group the meridian of T_1 is (up to conjugation and change of orientation) $\mu_1 = [x_1, t]$ and the meridian of T_2 is $\mu_2 = [x_2, s]$. $\square$

There is a standard procedure, due to Thurston [27], for constructing a symplectic structure on B for which G is a symplectic submanifold, and for which the tori T_1 and T_2 are Lagrangian. Since it is critical for us that these tori be Lagrangian, we give a proof of this fact for the convenience of the reader.

Lemma 4. *Let A denote an annulus $[0, 1] \times S^1$. Then there exists an area form α on A and a diffeomorphism $D : A \rightarrow A$ isotopic to a Dehn twist supported away from a neighborhood of the boundary so that $D^*(\alpha) = \alpha$.*

Proof. Let $g : [0, 1] \rightarrow [0, 2\pi]$ be a smooth non-decreasing function so that $g(x) = 0$ for $x < \frac{1}{3}$ and $g(x) = 2\pi$ for $x > \frac{2}{3}$. Define $\tilde{D} : [0, 1] \times \mathbb{R} \rightarrow [0, 1] \times \mathbb{R}$ by $\tilde{D}(x, y) = (x, y + g(x))$. Then $\tilde{D}^*(dx \wedge dy) = dx \wedge dy$. Since $\tilde{D}(x, y + 2\pi) = \tilde{D}(x, y)$, $\tilde{D}$ descends to a diffeomorphism of $[0, 1] \times S^1$ which represents a Dehn twist, preserves the standard area form, and is supported away from the boundary. $\square$

Corollary 5. *There exists a symplectic structure on B so that G is symplectic and T_1 and T_2 are Lagrangian. Moreover, B is minimal.*

Proof. By Lemma 4 we can find a Riemannian metric g on the surface G so that the Dehn twists D_1 and D_2 preserve the area form $\alpha(g)$ and are supported in small annular neighborhoods that miss the curve L . Let q_i $i = 1, 2, 3$ denote the projections of $G \times [0, 1] \times [0, 1]$. Then $\omega = q_1^*(\alpha(g)) + q_2^*(dt) \wedge q_3^*(ds)$ is a symplectic form on $G \times [0, 1] \times [0, 1]$ for which the slices $G \times \{(t, s)\}$ are symplectic and $K \times \{0\} \times I$ and $L \times I \times \{0\}$ are Lagrangian. Since $(D_2 D_1)^*(\alpha(g)) = \alpha(g)$ and since dt and ds descend to S^1 , ω descends to a well defined symplectic form on B for which G is symplectic and T_1 and T_2 are Lagrangian.

To see that B is minimal, it suffices to observe that the universal cover of B is contractible, and so $\pi_2(B) = 0$, since this implies that there are no spheres of self-intersection -1 . $\square$

Rescale the symplectic form on Matsumoto's manifold W if necessary so that the symplectic fiber $F = F_p \subset W$ is symplectomorphic to the symplectic surface $G \subset B$. Fix a symplectomorphism which takes the generators a_1, b_1, a_2, b_2 of $\pi_1(F)$ to $x_1, y_1, x_2, y_2 \in \pi_1(G)$ respectively.

Then form the symplectic sum $R = W \#_{F,G} B$ of W and B along F and G . As explained in Lemma 1, the fundamental group of R is obtained from $\pi_1(B)$ by setting $x_1 = x_2^{-1}$, $y_1 = y_2^{-1}$, and requiring x_1 and y_1 to commute. Thus:

$$\begin{aligned}\pi_1(R) &= \langle x_1, y_1, t \mid [x_1, y_1], [x_1, y_1][x_1^{-1}, y_1^{-1}], [t, x_1], [y_1^{-1}, t]x_1^{-1}, [x_1, t]y_1, [t, y_1^{-1}] \rangle \oplus \mathbb{Z}s \\ &= \mathbb{Z}t \oplus \mathbb{Z}s.\end{aligned}$$

In fact, the fourth and sixth relations force $x_1 = 1$, and then the fifth relation forces $y_1 = 1$.

The complement of $T_1 \cup T_2$ in R is the sum of W with $B - (T_1 \cup T_2)$ along F and G . Taking the presentation of $\pi_1(B - (T_1 \cup T_2))$ of Proposition 3, setting $x_1 = x_2^{-1}$, $y_1 = y_2^{-1}$, and requiring x_1 and y_1 to commute, one sees that $[t, y_1^{-1}] = 1$ and $[y_1^{-1}, t]x_1^{-1} = 1$. This forces $x_1 = 1$. Then $[x_1, t]y_1 = 1$ forces $y_1 = 1$. The presentation thus reduces to

$$\pi_1(R - (T_1 \cup T_2)) = \mathbb{Z}t \oplus \mathbb{Z}s.$$

In particular μ_1 and μ_2 are trivial in $\pi_1(R - (T_1 \cup T_2))$.

Gompf's symplectic sum theorem [12] guarantees that a symplectic structure can be found on R for which the tori T_1 and T_2 remain Lagrangian. Notice that each T_i represents a non-zero class in $H_2(R; \mathbb{R})$. In fact, the union of a small D^2 transverse to T_1 and a nullhomotopy of μ_1 in $R - (T_1 \cup T_2)$ gives a (singular) 2-sphere intersecting T_1 transversally in one point. This shows that T_1 represents a non-zero class in $H_2(R; \mathbb{R})$ and $H_2(R - T_2; \mathbb{R})$. Similarly T_2 represents a non-zero class in $H_2(R; \mathbb{R})$ and $H_2(R - T_1; \mathbb{R})$. The classes $[T_1]$ and $[T_2]$ are not multiples of each other since the singular 2-sphere dual to T_1 misses T_2 . It follows [12, Lemma 1.6] that the symplectic form on R can be perturbed by an arbitrarily small 2-form so that T_1 and T_2 become symplectic.

The Euler characteristic of R is computed using the formula $e(A \#_H B) = e(A) + e(B) - 2e(H)$ valid for any symplectic sum of 4-manifolds. From this formula we first see that $e(R) = e(W) + e(B) + 4 = 4 + 0 + 4 = 8$. Novikov additivity computes the signature $\sigma(R) = \sigma(W) + \sigma(B) = -4$. We will show in Theorem 11 below that R is minimal.

We summarize the properties of the manifold R in the following theorem

Theorem 6. *There exists a closed minimal symplectic 4-manifold R with Euler characteristic $e(R) = 8$ and signature $\sigma(R) = -4$ containing a disjoint pair of Lagrangian, homologically essential tori T_1, T_2 with trivial normal bundles so that $\pi_1(R) = \mathbb{Z}t \oplus \mathbb{Z}s$ and such that the homomorphism induced by the inclusion $\pi_1(R - (T_1 \cup T_2)) \rightarrow \pi_1(R)$ is an isomorphism.*

Moreover, the meridians μ_1 and μ_2 to the tori T_1 and T_2 are nullhomotopic in $R - (T_1 \cup T_2)$. The homomorphism induced by inclusion $\pi_1(T_1) \rightarrow \pi_1(R)$ takes one generator to s and the other to the identity and the homomorphism $\pi_1(T_2) \rightarrow \pi_1(R)$ takes one generator to t and the other to the identity. The symplectic form can be perturbed slightly so that one or both of the tori T_i become symplectic. $\square$

4. MODIFYING R BY LUTTINGER SURGERY

We can use Luttinger surgery [15] to modify R in a neighborhood of the Lagrangian torus T_1 (and T_2).

This symplectic construction is carefully explained in [1]. We recall the relevant details for the convenience of the reader.

An oriented Lagrangian torus T in a symplectic 4-manifold M has a tubular neighborhood symplectomorphic to a neighborhood of the zero section in its cotangent bundle by the Darboux theorem. Thus if x, y are oriented coordinates on T (i.e. we fix a universal covering $\mathbb{R}^2 \rightarrow T$) then dx, dy trivialize the cotangent bundle of T , and thereby one obtains a framing $T^2 \times D^2 \rightarrow M$ of a tubular neighborhood of T in M called the *Lagrangian framing*. As observed in [8], this framing can be described by the condition that $T^2 \times \{x\} \subset M$ is Lagrangian for all $x \in D^2$.

Luttinger showed that the manifold obtained by removing the tubular neighborhood $T \times D^2$ from M and regluing using an appropriate orientation preserving diffeomorphism $\psi : T \times S^1 \rightarrow T \times S^1$ yields a new manifold which admits a symplectic structure which agrees with the given symplectic structure on $M - T \times D^2$. To be precise, if we denote the generators of $H_1(T)$ (as well as their push offs into $H_1(T \times S^1)$)

using the Lagrangian framing by α, β , and the meridian of T by μ , Luttinger surgery yields a symplectic manifold if

$$\psi_*(\alpha) = \alpha, \psi_*(\beta) = \beta, \text{ and } \psi_*(\mu) = a\alpha + b\beta + \mu.$$

Let $k = \gcd(a, b)$ and set γ to be the embedded curve $\gamma = \frac{1}{k}(a\alpha + b\beta)$, we denote the resulting symplectic manifold by $M(\gamma, k)$. Notice that (identifying $\pi_1(T \times S^1)$ with $H_1(T \times S^1)$ and writing multiplicatively)

$$\pi_1(M(\gamma, k)) = \pi_1(M - T)/N(\gamma^k \mu).$$

This is because gluing in $T \times D^2$ can be accomplished by gluing in one 2 handle attached along $\gamma^k \mu$, then gluing in two 3-handles and one 4-handle.

The manifold $M(\gamma, k)$ is called $1/k$ Luttinger surgery on T along γ (the terminology comes from the observation that locally M is obtained by doing a $1/k$ Dehn surgery and crossing with S^1). Notice that $e(M(\gamma, k)) = e(M)$, and Novkov additivity shows that $\sigma(M(\gamma, k)) = \sigma(M)$.

We apply Luttinger surgery to $T_1 \subset R$. Take $\gamma = s$ and $k = 1$, and denote the resulting manifold by $R(s, 1)$ by P . Then

$$\pi_1(P) = \pi_1(R - T_1)/N(s\mu_1) = \mathbb{Z}t,$$

and $e(P) = 8$ and $\sigma(P) = -4$. Similarly

$$\pi_1(P - T_2) = \pi_1(R - (T_1 \cup T_2))/N(s\mu_1) = \mathbb{Z}t$$

and the meridian μ_2 to T_2 in $P - T_2$ is trivial, since it is trivial in $\pi_1(R - (T_1 \cup T_2))$.

Remark. In the preceding paragraph we did not mention the Lagrangian framing, although the reader may check that there is a curve on T_1 whose Lagrangian push off is $s \in \pi_1(R - (T_1 \cup T_2))$. An alternative construction of the manifold R is given in Section 5 in which it is obvious that the Lagrangian framing is obtained by pushing the curve K off itself in the fiber G (and taking the product with S^1).

However, note that if we take an *arbitrary* framing to push curves off T_i , the resulting push offs will differ from the Lagrangian framing push offs by some power of μ_1 . Since $\mu_1 = 1$ in $\pi_1(R - (T_1 \cup T_2))$, the fundamental group calculation is the same: s is killed, leaving t .

We will prove that P is minimal in Theorem 11 below.

We summarize these facts in the following theorem. To simplify the statement we denote T_2 simply by T .

Theorem 7. *There exists a closed minimal symplectic 4-manifold P with fundamental group $\mathbb{Z}$, Euler characteristic $e(P) = 8$ and signature $\sigma(P) = -4$ which contains a Lagrangian (or symplectic) homologically essential torus T with trivial normal bundle such that*

- (1) *The induced homomorphism $\pi_1(P - T) \rightarrow \pi_1(P)$ is an isomorphism,*
- (2) *The meridian of T is trivial in $\pi_1(P - T)$, and*
- (3) *the induced homomorphism $\pi_1(T) \rightarrow \pi_1(P)$ takes one symplectic generator to the generator of $\pi_1(P)$ and maps the other generator to the identity.*

□

Our interest in this manifold is two fold. First, it is the smallest known (to us) symplectic 4-manifold with fundamental group $\mathbb{Z}$, where we measure the size using the Euler characteristic (or equivalently the second Betti number). (Constructions of symplectic manifolds with fundamental group $\mathbb{Z}$ can be found in the literature, e.g. [18], [12], [24].) The other reason is that it can be used as a smaller replacement for the elliptic surface $E(1)$ typically used to control fundamental groups of symplectic 4-manifolds. We will illustrate this in subsequent sections, using the following theorem.

Theorem 8. *Let M be a symplectic 4-manifold containing a symplectic torus T' with trivial normal bundle such that $x, y \in \pi_1(M)$ represent the images of the two generators of $\pi_1(T')$. Then the symplectic sum of P and M along T and T' , $P \#_T M$, admits a symplectic structure (which agrees with that of P and M away from T, T') symplectic and satisfies*

$$e(P \#_T M) = e(M) + 8, \sigma(P \#_T M) = \sigma(M) - 4, \text{ and } \pi_1(P \#_T M) = \pi_1(M)/N(x)$$

where $N(x)$ denotes the normal subgroup of $\pi_1(M)$ generated by x . $\square$

In a different direction, we can perform Luttinger surgery on both Lagrangian tori T_1 and T_2 . Suppose we are given embedded curves $\gamma_1 \subset T_1$ representing $p_1 y_1 + q_1 s$ and $\gamma_2 \subset T_2$ representing $p_2 y_2 + q_2 t$, where p_1, q_1 are relatively prime integers and p_2, q_2 are relatively prime integers. Suppose further that a pair of integers k_1, k_2 are given.

Let $Q = Q(k_1, \gamma_1; k_2, \gamma_2)$ denote the symplectic 4-manifold obtained by performing Luttinger surgery so that $\gamma_1^{k_1} \mu_1$ and $\gamma_2^{k_2} \mu_2$ bound discs after regluing. We will prove that $Q(k_1, s; k_2, p y_2 + t)$ is minimal in Theorem 11 below.

Using Freedman's theorem [11] and Taubes's theorem [26] that minimal symplectic manifolds do not contain -1 spheres we immediately conclude the following.

Theorem 9. *The symplectic manifold $Q(k_1, p_1 y_1 + q_1 s; k_2, p_2 y_2 + q_2 t)$ has Euler characteristic 8 and signature -4 , and fundamental group $\mathbb{Z}/(k_1 q_1) \oplus \mathbb{Z}/(k_2 q_2)$. It is minimal when $p_1 = 0, q_1 = 1$ and $q_2 = 1$.*

In particular, $Q(1, p_1 y_1 + s; 1, p_2 y_2 + t)$ is simply connected and hence is homeomorphic to $\mathbb{C}P^2 \# 5\overline{\mathbb{C}P}^2$. The manifolds $Q(1, s; 1, p y_2 + t)$ (indexed by $p \in \mathbb{Z}$) are minimal, hence not diffeomorphic to $\mathbb{C}P^2 \# 5\overline{\mathbb{C}P}^2$. $\square$

Looking from the inside out, we can define $M = Q(1, s; 1, t)$. Then M is a minimal symplectic manifold homeomorphic to $\mathbb{C}P^2 \# 5\overline{\mathbb{C}P}^2$ which contains a (nullhomologous) Lagrangian torus T_2 . Moreover, for each integer p , the manifold $M(p) = Q(1, s; 1, p y_2 + t)$ is another minimal symplectic manifold homeomorphic to $\mathbb{C}P^2 \# 5\overline{\mathbb{C}P}^2$, obtained from M by performing Luttinger surgery on T_2 . It seems reasonable to conjecture that these are all smoothly distinct.

The construction of manifold homeomorphic but not diffeomorphic to $\mathbb{C}P^2 \# k\overline{\mathbb{C}P}^2$ s for $k \leq 9$ began with Donaldson's seminal example [7] that the Dolgachev surface $E(1)_{2,3}$ is not diffeomorphic to $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P}^2$. In 1989, Kotschick [14] proved that the Barlow surface is homeomorphic but not diffeomorphic to $\mathbb{C}P^2 \# 8\overline{\mathbb{C}P}^2$. In 2004 J. Park [21] constructed the first exotic smooth structure on $\mathbb{C}P^2 \# 7\overline{\mathbb{C}P}^2$. Since then Park's results have been expanded upon in [19, 25, 9, 23], producing infinite families of smooth 4-manifolds homeomorphic but not diffeomorphic to $\mathbb{C}P^2 \# k\overline{\mathbb{C}P}^2$ for $k = 5, 6, 7, 8$. The $n = 5$ examples are not symplectic.

In [2], Akhmedov produced the first example of an symplectic 4-manifold homeomorphic to but not diffeomorphic to $\mathbb{C}P^2 \# 5\overline{\mathbb{C}P}^2$. He uses a different construction from ours, although the manifold W is a key ingredient in his construction, as in ours. Very recently Akhmedov, Baldridge, and Park [3, 5] have constructed minimal symplectic manifolds homeomorphic to $\mathbb{C}P^2 \# 3\overline{\mathbb{C}P}^2$.

There is also an interesting literature on the construction of manifolds homeomorphic but not diffeomorphic to $3\mathbb{C}P^2 \# n\overline{\mathbb{C}P}^2$ for small n . D. Park [20] has constructed many such manifolds and Akhmedov [2] article contains the construction of a minimal symplectic manifold homeomorphic to $3\mathbb{C}P^2 \# 7\overline{\mathbb{C}P}^2$. In their recent work Akhmedov, Baldridge, and Park [3, 5] have constructed minimal symplectic manifolds homeomorphic to $3\mathbb{C}P^2 \# 5\overline{\mathbb{C}P}^2$.

Note that the smallest previously known symplectic 4-manifolds with cyclic fundamental group have $e \geq 10$. There are smooth manifolds with cyclic fundamental group and $e = 2$, and any closed 4-manifold with cyclic fundamental group has $e \geq 2$. In any case, any symplectic 4-manifold with cyclic fundamental group has $e + \sigma \geq 4$.

5. MINIMALITY

In this section we prove that the manifolds constructed above are minimal. The key result is the following theorem.

Theorem 10. *Let $T^4 = (S^1 \times S^1) \times (S^1 \times S^1)$ be given the product symplectic form. Let $T_1 = S^1 \times \{a\} \times S^1 \times \{a\}$, and $T_2 = S^1 \times \{b\} \times \{b\} \times S^1$ where $a \neq b$. Denote by $x, y, z, w \in \pi_1(T^4)$ the generators given by the coordinate circles.*

Then the symplectic manifold M obtained from $1/k_1$ Luttinger surgery on T_1 along x and $1/k_2$ Luttinger surgery on T_2 along w is minimal.

Proof. Notice first that the Lagrangian framings for the tori T_1 and T_2 are obvious: the parallel Lagrangian tori are obtained by varying the points a and b .

Let M denote the symplectic manifold obtained by $1/k_1$ Luttinger surgery on T_1 along x and $1/k_2$ Luttinger surgery on T_2 along w .

It is straightforward to check that $1/k_1$ surgery on T_1 along x transforms $T^4 = (T^3) \times S^1$ into $N \times S^1$, where N is the 3-manifold that fibers over S^1 with monodromy $D_x^{k_1} : T^2 \rightarrow T^2$, where D_x denotes the Dehn twist on T^2 along the first coordinate. This is explained in [1, pg. 189].

View $N \times S^1$ as a trivial circle bundle over N . Then the fiber is represented by the curve w and T_2 is the restriction of this circle bundle to $x \times \{(b, b)\}$ in N . Luttinger surgery on T_2 along w corresponds to twisting the bundle over x , i.e. replacing $N \times S^1 \rightarrow N$ by the S^1 bundle $M \rightarrow N$ with first Chern class equal to k_2 times the Poincaré dual of x in N . Details can be found in [4]. In any case one can check directly from the formula that M has a free circle action which coincides with the action on $N \times S^1$ away from T_2 .

Thus M is an S^1 bundle over a fibered 3-manifold N with fiber a torus. It follows from the long exact sequence of homotopy groups that $\pi_2(M) = 0$, and hence M contains no essential 2-spheres. In particular, M is minimal. $\square$

If $S : T^4 \rightarrow T^4$ is a symplectic diffeomorphism then obviously the manifold obtained by $1/k_1$ surgery on $S(T_1)$ along $S(x)$ and $1/k_2$ surgery on $S(T_2)$ along $S(w)$ is also minimal. So for example, the linear transformation $\tilde{S} : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ given by $\tilde{S}(x, y, z, w) = (x, y + pz, z, px + w)$ descends to a symplectic diffeomorphism of T^4 which leaves T^1 and T^2 invariant. Thus the manifold obtained by $1/k_1$ surgery on T_1 along x and $1/k_2$ surgery on T_2 along $x^p w$ is also minimal.

Given an integer k_1 , let $M_1(k_1)$ denote the symplectic manifold obtained from T^4 by two Luttinger surgeries: $1/1$ surgery on $T_0 = S^1 \times \{a\} \times \{a\} \times S^1$ along x and $1/k_1$ surgery on $T_1 = \{b\} \times S^1 \times \{b\} \times S^1$ along w . After suitably permuting coordinates, Theorem 10 implies that $M_1(k_1)$ is minimal. Note that T^4 contains a pair of (intersecting) symplectic tori $R_1 = \{c\} \times \{c\} \times S^1 \times S^1$ and $G_1 = S^1 \times S^1 \times \{d\} \times \{d\}$ disjoint from T_0 and T_1 ; these tori remain symplectic in $M_1(k_1)$.

Given a pair of integers p and k_2 , let $M_2(p, k_2)$ denote the symplectic manifold obtained from T^4 by two Luttinger surgeries: $1/1$ surgery on $T_3 = \{a\} \times S^1 \times \{a\} \times S^1$ along y and $1/k_2$ surgery on $T_2 = \{b\} \times S^1 \times S^1 \times \{b\}$ along $y^p z$. Theorem 10 implies that $M_2(p, k_2)$ is minimal. Again, T^4 contains a pair of symplectic tori $R_2 = \{c\} \times \{c\} \times S^1 \times S^1$ and $G_2 = S^1 \times S^1 \times \{d\} \times \{d\}$ disjoint from T_3 and T_2 ; these tori remain symplectic in $M_2(p, k_2)$.

The fiber sum of $M_1(k_1)$ and $M_2(p, k_2)$ along R_1 and R_2 is therefore a minimal symplectic manifold by Usher's theorem [28]. Denote this manifold by $M(k_1, p, k_2)$. Note that the two symplectic surfaces G_1 and G_2 can be matched up to produce a symplectic genus 2 surface $G \subset M(k_1, p, k_2)$ [12]. Taking the symplectic sum of $M(k_1, p, k_2)$ with the manifold W along the genus two surfaces G and F again produces a minimal symplectic manifold by Usher's theorem since the exceptional spheres in W all intersect the genus two surface F .

A straightforward examination of the constructions shows that this fiber sum $M(k_1, p, k_2) \#_G W$ is exactly the symplectic manifold denoted by $Q(k_1, \gamma_1; k_2, \gamma_2)$ in Section 4, where $\gamma_1 = s$ (so $p_1 = 0, q_1 = 1$ in the notation of Section 4) and $\gamma_2 = py_2 + t$ (so $p_2 = p, q_2 = 1$). In fact, note that the fiber sum of two copies of T^4 along R_1 and R_2 gives $G \times T^2$. The Luttinger surgeries along the tori T_0 and T_3 correspond exactly to replacing the trivial bundle $G \times S^1 \rightarrow S^1$ by the fiber bundle with monodromy the Dehn twists D_1 and D_2 . The remaining tori T_1 and T_2 coincide with the tori labelled T_1 and T_2 in Section 3 and one checks that the surgery coefficients are correct. We summarize these facts in the following theorem.

Theorem 11. *For any triple of integers k_1, k_2, p , the symplectic manifold $Q(k_1, s; k_2, py_2 + t)$ is minimal. Its fundamental group is given by*

$$\pi_1(Q(k_1, s; k_2, py_2 + t)) = \mathbb{Z}/k_1 \oplus \mathbb{Z}/k_2.$$

It has Euler characteristic equal to 8 and signature -4 .

In particular, the symplectic manifold $R = Q(0, s, 0, t)$ with fundamental group $\mathbb{Z}^2$ of Theorem 6 is minimal, the manifold $P = Q(1, s, 0, t)$ with infinite cyclic fundamental group of Theorem 7 is minimal and the simply connected manifolds $M(p) = Q(1, s; 1, py_2 + t)$ of Theorem 9 are minimal (and hence exotic) manifolds. $\square$

Remark. We have not tried to be as general as possible in Theorem 11. Presumably a more careful examination will reveal that $Q(k_1, \gamma_1; k_2, \gamma_2)$ is minimal for arbitrary γ_1 and γ_2 . We believe our approach has much potential: one can consider more complicated monodromies, symplectic sums of more than two copies of T^4 , and hence surfaces of higher genus and surgeries along more Lagrangian tori, and the use of other manifolds besides W .

6. MORE APPLICATIONS.

6.1. Free abelian groups. We first give an easy application of Theorem 8. Let M denote the 4-torus with its natural symplectic structure and $T' \subset M$ one of the coordinate symplectic tori.

Corollary 12. $P \#_T M$ is a symplectic 4-manifold with fundamental group $\mathbb{Z}^3$, Euler characteristic 8, and signature -4 . $\square$

The smallest previously known example of a symplectic 4-manifold with fundamental group $\mathbb{Z}^3$ has $e = 12$ and any such symplectic manifold must have $e \geq 3$ ([6]).

More generally, the technique of [6, Theorem 20] allows us to improve the construction of symplectic 4-manifolds with odd rank free abelian groups by taking the fiber sum of $Sym^2(F_n)$ (F_n a surface of genus n) with the manifold P , rather than the larger manifold K of [6, Lemma 18]. We refer the interested reader to loc.cit. for details of the proof of the following corollary.

Corollary 13. There exists a symplectic 4-manifold M with $\pi_1(M) = \mathbb{Z}^{2n-1}$ such that $e(M) = 11 - 5n + 2n^2$ and $\sigma(M) = -3 - n$. $\square$

This is an improvement over the upper bound $\min_{\pi_1(M)=\mathbb{Z}^{2n-1}} e(M) \leq 15 - 5n + 2n^2$ of [6, Theorem 20], but still far from the lower bound $6 - 7n + 2n^2 \leq \min_{\pi_1(M)=\mathbb{Z}^{2n-1}} e(M)$. Better constructions are needed to decrease the upper bound (or to increase the lower bound).

6.2. Arbitrary fundamental groups. As explained in Section 4 of [6], the existence of the symplectic manifold P and its symplectic torus T of Theorem 8 allows us to improve (by 30%) the main result of loc.cit. to the following theorem.

Theorem 14. Let G be a finitely presented group that has a presentation with g generators and r relations. Then there exists a symplectic 4-manifold M with $\pi_1 M \cong G$, Euler characteristic $e(M) = 12 + 8(g+r)$, and signature $\sigma(M) = -8 - 4(g+r)$. Moreover, M contains a symplectic torus that lies in a cusp neighborhood. $\square$

Proof. In the proof of [6, Theorem 6] a symplectic 4-manifold N is constructed whose fundamental group contains classes $s, t, \gamma_1, \dots, \gamma_{r+g}$ so that G is isomorphic to the quotient of $\pi_1(N)$ by these classes. Moreover, N contains symplectic tori $T_0, T_1, \dots, T_{g+r}$ so that the two generators of $\pi_1(T_0)$ represent s and t , and for $i \geq 1$ the two generators of $\pi_1(T_i)$ represent s and γ_i .

Let $E(1)$ denote the elliptic surface with $e(E(1)) = 12$, the fibration chosen to have (at least) two cusp fibers. Let F_1, F_2 be regular fibers with F_1 near one cusp and F_2 near another. Take the fiber sum of N with $E(1)$ along F_1 , and $g+r$ copies of the manifold P of Theorem 7 along the tori T_i , $i \geq 1$ in such a way that the γ_i are killed. Then a repeated application of Theorem 8 computes

$$\pi_1(M) = \pi_1(N)/N(s, t, \gamma_i) = G, \quad e(M) = 12 + 8(g+r), \quad \sigma(M) = -8 - 4(g+r).$$

The torus F_2 survives in M as a symplectic torus which lies in a cusp neighborhood. $\square$

An examination of the proof of Theorem 6 of [6] shows that Theorem 14 can be improved for certain presentations, namely, one can find M so that $e(M) = 12 + 8(g' + r)$ and $\sigma(M) = -8 - 4(g' + r)$, where g' is the number of generators which appear in some relation with negative exponent. Thus if G has a presentation with r relations in which every generator appears only with positive exponent in each relation, then there exists a symplectic M with $\pi_1(M) = G$, and $e(M) = 12 + 8r$, $\sigma(M) = -8 - 4r$. Moreover, using Usher's theorem [28] one sees that the manifolds constructed are minimal.

These manifolds can all be assumed to contain a symplectic torus in a cusp neighborhood, since we use one copy of $E(1)$ to get the construction started. Thus the geography results of J. Park [22] can be improved to find a larger region of the (c_1^2, χ_h) plane for which to each pair of integers in that region one can find infinitely many non-diffeomorphic, homeomorphic minimal symplectic manifolds with fundamental group G .

Another (decidedly minor) improvement concerns groups of the form $G \times \mathbb{Z}$: for a presentation of G as above there exists a symplectic manifold M with $\pi_1(M) = G \times \mathbb{Z}$, $e(M) = 8(g' + r + 1)$, and $\sigma(M) = -4(g' + r + 1)$. The reason is that one step in the proof of [6, Theorem 6] consists of taking a symplectic sum with $E(1)$ to kill two generators (t and s). But to get $G \times \mathbb{Z}$ it suffices to kill t , for which the manifold P of Theorem 8 can be used instead of $E(1)$. (Of course, if G is the fundamental group of a 3-manifold Y that fibers over S^1 , then $Y \times S^1$ is a symplectic 4-manifold with fundamental group $G \times \mathbb{Z}$ and Euler characteristic zero.)

Suppose M is a symplectic 4-manifold containing a symplectic torus T with trivial normal bundle so that $\pi_1(M - T) = 1$ or $\mathbb{Z}$ and so that pushing T into $M - T$ induces a surjection $\pi_1(T) \rightarrow \pi_1(M - T)$. It is easy to see that such an M must have $b^+ > 1$ and $b^- > 0$, and hence if M is simply connected $e(M) \geq 6$. If $\pi_1(M) = \mathbb{Z}$ such an M must have $e(M) \geq 3$. The manifold P has $e(P) = 8$. Further improvements in the geography problem for symplectic manifolds will be obtained if such an M is found with $e(M) < 8$. The search for such a manifold is a promising direction for future study.

6.3. Other small simply connected manifolds. Our manifolds can be used to produce more interesting examples of small simply connected symplectic manifolds. We consider the case when $b^+ = 3$.

As a warm up, consider the fiber sum of two copies of P along T (see Theorem 7). If we form the fiber sum along a symplectic diffeomorphism that interchanges (up to sign) the two generators of $\pi_1(T)$, the resulting manifold is a minimal simply connected symplectic manifold homeomorphic to $3\mathbb{CP}^2 \# 11\overline{\mathbb{CP}}^2$. One can glue using other maps to obtain manifolds with cyclic fundamental group with $e = 16$ and $\sigma = -8$, reminiscent of the way one constructs lens spaces from two solid tori. In particular there are infinitely many (isotopy classes of) gluings which produce simply connected examples. We do not know if these give distinct diffeomorphism types.

For a smaller example, we start with a symplectic genus 2 surface H of square zero in $T^4 \# 2\overline{\mathbb{CP}}^2$ constructed by symplectically resolving $T^2 \times \{(a, a)\} \cup \{(a, a)\} \times T^2$ and blowing up twice. Thus the homomorphism induced on fundamental groups by the inclusion $H \subset T^4 \# 2\overline{\mathbb{CP}}^2$ takes the four standard generators of the genus 2 surface group to the four coordinate generators of $\pi_1(T^4 \# 2\overline{\mathbb{CP}}^2) = \mathbb{Z}^4$. Since H intersects each exceptional sphere in one point, the meridian of H in $T^4 \# 2\overline{\mathbb{CP}}^2$ is nullhomotopic. Hence an argument exactly like the proof of Lemma 1 shows that the symplectic sum of a manifold X with $T^4 \# 2\overline{\mathbb{CP}}^2$ along a square zero genus 2 surface has fundamental group obtained from that of X by requiring the four generators of the genus 2 surface in X to commute.

Consider a parallel copy F' of the symplectic surface F in the manifold $W = (T^2 \times S^2) \# 4\overline{\mathbb{CP}}^2$. Then the sum W' of W and $T^4 \# 2\overline{\mathbb{CP}}^2$ along F' and H is symplectic and minimal, using Usher's theorem. Moreover, $\pi_1((W - F) \#_{F'} (T^4 \# 2\overline{\mathbb{CP}}^2)) = \mathbb{Z}^2$, and a quick check shows that the statement of Lemma 1 holds with W' replacing W . The difference between the two is that $e(W') = 10$ and $\sigma(W') = -6$.

Taking the fiber sum R' of W' (rather than W) with B , where B is the manifold constructed in Section 3, and performing Luttinger surgery gives another family $Q'(k_1, \gamma_1; k_2, \gamma_2)$ with $e = 14$ and $\sigma = -6$. As in Theorem 9 the manifolds $M'(p) = Q'(1, s, py_2 + t)$ are a family of minimal (hence exotic) simply connected

symplectic manifolds homeomorphic to $3\mathbb{CP}^2 \# 9\overline{\mathbb{CP}}^2$. We remark that $M'(p)$ is obtained from $M'(0)$ by $1/p$ Luttinger surgery.

6.4. Some comments on Seiberg-Witten calculations. Fix k and consider the family $M(k, p) = Q(k_1, s; 1, py_2 + t)$ of minimal symplectic manifolds with cyclic fundamental group $\mathbb{Z}/k$. Then $M(k, p)$ can also be described as $1/p$ Luttinger surgery on $C(k, 0)$ along y_2 . The usual way ([10]) to distinguish such families of manifolds is by their Seiberg-Witten invariants, as follows.

The torus T_2 is nullhomologous in $M(k, p)$ for all p and so the Morgan-Mrowka-Szabo formula [17] for the Seiberg-Witten invariants of $M(k, p)$ (see also [10, Theorem 5.3]) states that

$$SW_{M(k,p)}(\kappa_p) = SW_{M(k,0)}(\kappa_0) + p \sum_i SW_{X(k)}(\kappa_X + i[T])$$

where $X(k)$ is the manifold obtained from $M(k, 0) - nbd(T_2)$ by gluing on $T^2 \times D^2$ in such a way that the curve y_2 bounds the meridian, and κ_p, κ_X are homology classes that correspond (i.e. agree in $H_2(M(k, 0), T_2)$).

However, the manifold $X(k)$ is not obtained by Luttinger surgery, and need not be symplectic. Hence we do not know if $\sum_i SW_{X(k)}(\kappa_X + i[T])$ is non-zero and it does not immediately follow that the manifolds $M(k, p)$ are distinct. In fact, when $k = 1$ (the case of exotic $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$) a theorem of Stipicz and Szabo [25, Proposition 4.3] implies that $\sum_i SW_{X(1)}(\kappa_X + i[T]) = 0$. In particular, these Seiberg-Witten invariants cannot distinguish the manifolds $M(1, p)$.

REFERENCES

- [1] D. Auroux, S.K. Donaldson, and L. Katzarkov, *Luttinger surgery along Lagrangian tori and non-isotopy for singular symplectic plane curves*. Math. Ann. 326 (2003), no. 1, 185–203.
- [2] A. Akhmedov, *Exotic smooth structures on $3\mathbb{CP}^2\#7\overline{\mathbb{CP}}^2$* . preprint (2006) math.GT/0612130
- [3] A. Akhmedov, B. Doug Park, *Exotic Smooth Structures on Small 4-Manifolds*. preprint (2007).
- [4] S. Baldridge, *New symplectic 4-manifolds with $b_+ = 1$* . Mathematische Annalen 333 (2005) 633–643.
- [5] S. Baldridge, *An exotic symplectic structure on $\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$ and on $3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$* . Preprint (2007).
- [6] S. Baldridge and P. Kirk, *On symplectic 4-manifolds with prescribed fundamental group*. To appear in Commentarii Mathematici Helvetici.
- [7] S. Donaldson, *Irrationality and the h-cobordism conjecture*. J. Differential Geom. 26 (1987), no. 1, 141–168.
- [8] R. Fintushel and R. Stern, *Invariants for Lagrangian tori*. Geom. Topol. 8 (2004), 947–968
- [9] R. Fintushel and R. Stern, *Double node neighborhoods and families of simply connected 4-manifolds with $b^+ = 1$* . J. Amer. Math. Soc. 19 (2006), no. 1, 171–180.
- [10] R. Fintushel and R. Stern, *Surgery on nullhomologous tori and simply connected 4-manifolds with $b^+ = 1$* . Preprint (2007) math.GT/0701044.
- [11] M. Freedman, *The topology of four-dimensional manifolds*. J. Differential Geom. 17 (1982), no. 3, 357–453.
- [12] R. Gompf, *A new construction of symplectic manifolds*. Ann. of Math. (2) 142 (1995), no. 3, 527–595.
- [13] R. Gompf and A. Stipsitz, *4-manifolds and Kirby calculus*. Graduate Studies in Mathematics, 20. American Mathematical Society, Providence, RI, 1999.
- [14] D. Kotschick, *On manifolds homeomorphic to $\mathbb{CP}^2\#8\overline{\mathbb{CP}}^2$* . Invent. Math. 95 (1989), no. 3, 591–600.
- [15] K. M. Luttinger, *Lagrangian Tori in $\mathbb{R}^4$* . J. Diff. Geom. 52 (1999), 203–222.
- [16] Y. Matsumoto, *Lefschetz fibrations of genus two—a topological approach*. Topology and Teichmüller spaces (Katinkulta, 1995), 123–148, World Sci. Publ., River Edge, NJ, 1996.
- [17] J. Morgan, T. Mrowka, and Z. Szabo, *Product formulas along T^3 for Seiberg-Witten invariants*. Math. Res. Lett. 4 (1997), no. 6, 915–929.
- [18] B. Ozbagci and A. Stipsitz, *Noncomplex smooth 4-manifolds with genus 2 Lefschetz fibrations*. Proc. Amer. Math. Soc. 128 (2000), no. 10, 3125–3128.
- [19] P. Ozsváth and Z. Szabó, *On Park’s exotic smooth four-manifolds*, ‘Geometry and topology of manifolds’, 253–260, Fields Inst. Commun., 47, Amer. Math. Soc., Providence, RI, 2005.
- [20] B. Doug Park, *Exotic smooth structures on $3\mathbb{CP}^2\#n\overline{\mathbb{CP}}^2$* . Proc. Amer. Math. Soc. 128 (2000), no. 10, 3057–3065.
- [21] Jongil Park, *Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* . Invent. Math. 159 (2005), no. 3, 657–667.
- [22] Jongil Park, *The geography of symplectic 4-manifolds with arbitrary fundamental group*. Preprint (2006).
- [23] Jongil Park, A. Stipsicz, and Z. Szabo, *Exotic smooth structures on $\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$* . Math. Res. Lett. 12 (2005), no. 5–6, 701–712.
- [24] I. Smith, *Symplectic geometry of Lefschetz fibrations*. Dissertation, Oxford 1998.
- [25] A. Stipsicz and Z. Szabó, *An exotic smooth structure on $\mathbb{CP}^2\#6\overline{\mathbb{CP}}^2$* , Geom. Topol. 9 (2005), 813–832.
- [26] C. Taubes, .
- [27] W. Thurston, *Some simple examples of symplectic manifolds*. Proc. Amer. Math. Soc. 55 (1976), 467–468.
- [28] M. Usher, *Minimality and symplectic sums*. To appear in Internat. Math. Res. Not. preprint (2006) math.SG/0606543

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY

BATON ROUGE, LA 70817

E-mail address: sbaldr@math.lsu.edu

MATHEMATICS DEPARTMENT, INDIANA UNIVERSITY

BLOOMINGTON, IN 47405

E-mail address: pkirk@indiana.edu